

Thurston's norm revisited

Igor V. Nikolaev *

Abstract

We study the Thurston norm on the second homology of a 3-manifold M , which is the surface bundle over the circle with a pseudo-Anosov monodromy. A novelty of our approach consists in the application of the C^* -algebras to a problem in topology. Namely, one associates to M a C^* -algebra, whose K -theory gives rise to an algebraic number field K . It is shown, that the trace function on the ring of integers of K induces a norm on the second homology of M . The norm coincides with the Thurston norm on the second homology of M .

Key words and phrases: operator algebras, 3-manifolds

AMS (MOS) Subj. Class.: 19K, 46L, 57M.

1 Introduction

In 1986, W. P. Thurston discovered the fundamental measure of complexity of a 3-dimensional manifold, M . The measure is a norm on the second homology of M , which assigns the non-negative integers to the elements of the group $H_2(M) = H_2(M; \mathbb{Z})$. If $z \in H_2(M)$, then a number $N(z)$ is attached, such that $N(z) = \min_X \{-\chi(X) \mid X \neq S^2 \text{ is compact surface representing class } z\}$, where χ is the Euler number of X . The norm is called a *Thurston norm*. The function N is linear on the $H_2(M)$ and extends to the real homology group $H_2(M; \mathbb{R})$ as a pseudo-norm (Thurston [10]). The Thurston norm is an important homotopy invariant of the manifold M , which can be viewed as a generalization of the genus of a knot.

It is interesting that the abelian groups with a norm arise in the context of the AF -algebras (the operator algebras, see Effros [2]). Namely, let $\mathbb{Z}^k \xrightarrow{A} \mathbb{Z}^k \xrightarrow{A} \mathbb{Z}^k \xrightarrow{A} \dots$, be a stationary dimension group. (We refer the reader to the section 2 for a definition.) By λ_A one understands the Perron-Frobenius eigenvalue of the positive integral matrix A . Let $K = \mathbb{Q}(\lambda_A)$ be an algebraic

*Partially supported by NSERC.

number field of the degree k , obtained as an extension of the rationals by the algebraic number λ_A . The field K is known to be an important invariant of the stationary dimension group. Namely, the triple (K, α, I) , where α is an embedding of the field K and I is the equivalence class of ideals in the ring of integers of K , is a complete Morita invariant of the stationary dimension group (Bratteli, Jørgensen, Kim & Roush [1], Effros [2], Handelmann [6]). Denote by O_K the ring of integers of the field K and fix an integral basis $\omega_1, \dots, \omega_k$ in O_K . Note that $O_K \cong \mathbb{Z}^k$ by the Gauss isomorphism. It is well known that the multiplication by $\alpha \in O_K$ induces a linear operator on the vector space O_K :

$$\begin{pmatrix} \alpha\omega_1 \\ \vdots \\ \alpha\omega_k \end{pmatrix} = \begin{pmatrix} a_{11} & \dots & a_{1k} \\ \vdots & & \vdots \\ a_{k1} & \dots & a_{kk} \end{pmatrix} \begin{pmatrix} \omega_1 \\ \vdots \\ \omega_k \end{pmatrix},$$

where a_{ij} are the rational integers. Define a function $\mathcal{N} : O_K \rightarrow \mathbb{Z}$ by the formula $\alpha \mapsto a_{11} + \dots + a_{kk}$, where $\alpha \in O_K$. It is not hard to verify, that $\mathcal{N}$ is a linear function, which is independent of the choice of the integral basis in O_K (Weyl [13]). Note that the pre-image $\mathcal{N}^{-1}(\mathbb{Z}^+)$ of the semi-group $\mathbb{Z}^+ = \{0, 1, 2, \dots\}$ is a cone $C \subset \mathbb{Z}^k$.

The aim of this note is to show that the algebraic norm $\mathcal{N}$ and the Thurston norm N are related. Namely, let $\varphi : X \rightarrow X$ be a pseudo-Anosov diffeomorphism of the compact surface X of the genus g and let $\mathcal{F}$ be the φ -invariant foliation on X (Thurston [11]). Consider the mapping torus of φ , i.e. a 3-dimensional manifold $M = \{X \times [0, 1] \mid (x, 0) \mapsto (\varphi(x), 1), \ x \in X\}$. (The reader can recognize M to be the surface bundle over the circle with a monodromy φ .) Let us construct the crossed product C^* -algebra $A_\varphi = C(X) \rtimes_\varphi \mathbb{Z}$, where $C(X)$ is the C^* -algebra of the continuous complex-valued functions on the surface X . It can be shown, that the K_0 -group of A_φ is a stationary dimension group. Define a map $\mathcal{N} : H_2(M) \rightarrow \mathbb{Z}$ using a natural isomorphism $O_K \cong H_1(X, \text{Sing } \mathcal{F}; \mathbb{Z}) \cong H_2(M)$, where $\text{Sing } \mathcal{F}$ is the set of singular points of the foliation $\mathcal{F}$. Our main result is the following theorem.

Theorem 1 *For every surface bundle $M \rightarrow S^1$ with a pseudo-Anosov monodromy φ , the following is true: (i) the preimage $\mathcal{N}^{-1}(\mathbb{Z}^+)$ of the semi-group $\mathbb{Z}^+ = \{0, 1, 2, \dots\}$ is a cone $C \subset H_2(M)$ and (ii) the norm $\mathcal{N}$ coincides with the Thurston norm N on the cone C .*

The structure of the note is the following. In section 2, the notation is introduced. Theorem 1 is proved in section 3.

2 Notation

This section is a brief introduction to the dimension groups, the algebraic number fields and the Thurston norm on the 3-dimensional manifolds. We refer

the reader to O. Bratteli, P. E. T. Jørgensen, K. H. Kim & F. Roush, ([1]), M. Rørdam, F. Larsen & N. Laustsen ([9]), H. Weyl ([13]) and W. Thurston ([10]) for a complete account.

2.1 The dimension group

By the C^* -algebra one understands a noncommutative Banach algebra with an involution ([9]). Namely, a C^* -algebra A is an algebra over $\mathbb{C}$ with the norm $a \mapsto \|a\|$ and an involution $a \mapsto a^*, a \in A$, such that A is complete with respect to the norm, and such that $\|ab\| \leq \|a\| \|b\|$ and $\|a^*a\| = \|a\|^2$ for every $a, b \in A$. If A is commutative, then the Gelfand theorem says that A is isometrically $*$ -isomorphic to the C^* -algebra $C_0(X)$ of the continuous complex-valued functions on a locally compact Hausdorff space X . For otherwise, A represents a noncommutative topological space.

2.1.1 The ordered abelian groups

Given a C^* -algebra, A , consider a new C^* -algebra $M_n(A)$, i.e. the matrix algebra over A . There exists a remarkable semi-group, A^+ , connected to the set of projections in the algebra $M_\infty = \cup_{n=1}^\infty M_n(A)$. Namely, the projections $p, q \in M_\infty(A)$ are the Murray-von Neumann equivalent $p \sim q$, if they can be presented as $p = v^*v$ and $q = vv^*$ for an element $v \in M_\infty(A)$. An equivalence class of the projections is denoted by $[p]$. The semi-group A^+ is defined to be the set of all equivalence classes of projections in $M_\infty(A)$ with the binary operation $[p] + [q] = [p \oplus q]$. The Grothendieck completion of A^+ to an abelian group is called a K_0 -group of A . The functor $A \rightarrow K_0(A)$ maps the unital C^* -algebras into a category of the abelian groups, so that the semi-group $A^+ \subset A$ corresponds to a positive cone $K_0^+ \subset K_0(A)$ and the unit element $1 \in A$ corresponds to an order unit $[1] \in K_0(A)$. The ordered abelian group $(K_0, K_0^+, [1])$ with the order unit is called a *dimension group* of the C^* -algebra A .

2.1.2 The AF-algebras

An *AF* (approximately finite-dimensional) C^* -algebra is defined to be a norm closure of an ascending sequence of the finite dimensional C^* -algebras M_n 's, where M_n is a C^* -algebra of the $n \times n$ matrices with the entries in $\mathbb{C}$. Here the index $n = (n_1, \dots, n_k)$ represents a *multi-matrix* C^* -algebra $M_n = M_{n_1} \oplus \dots \oplus M_{n_k}$. Let $M_1 \xrightarrow{\varphi_1} M_2 \xrightarrow{\varphi_2} \dots$, be a chain of the finite-dimensional C^* -algebras and their homomorphisms. A set-theoretic limit $A = \lim M_n$ has a natural algebraic structure given by the formula $a_m + b_k \rightarrow a + b$; here $a_m \rightarrow a, b_k \rightarrow b$ for the sequences $a_m \in M_m, b_k \in M_k$. The homeomorphisms of the above (multi-matrix) algebras admit a canonical description (Effros [2]). Suppose that $p, q \in \mathbb{N}$ and $k \in \mathbb{Z}^+$ are such numbers that $kq \leq p$. Let us define a homomorphism $\varphi : M_q \rightarrow M_p$ by the formula $a \mapsto a \oplus \dots \oplus a \oplus 0_h$, where $p = kq + h$. More

generally, if $q = (q_1, \dots, q_s), p = (p_1, \dots, p_r)$ are vectors in $\mathbb{N}^s, \mathbb{N}^r$, respectively, and $\Phi = (\phi_{kl})$ is a $r \times s$ matrix with the entries in $\mathbb{Z}^+$ such that $\Phi(q) \leq p$, then the homomorphism φ is defined by the formula:

$$\begin{aligned} a_1 \oplus \dots \oplus a_s &\longrightarrow \underbrace{(a_1 \oplus a_1 \oplus \dots)}_{\phi_{11}} \oplus \underbrace{(a_2 \oplus a_2 \oplus \dots)}_{\phi_{12}} \oplus \dots \oplus 0_{h_1} \\ &\oplus \underbrace{(a_1 \oplus a_1 \oplus \dots)}_{\phi_{21}} \oplus \underbrace{(a_2 \oplus a_2 \oplus \dots)}_{\phi_{22}} \oplus \dots \oplus 0_{h_2} \oplus \dots \end{aligned}$$

where $\Phi(q) + h = p$. We say that φ is a *canonical homomorphism* between M_p and M_q . Any homomorphism $\varphi : M_q \rightarrow M_p$ can be rendered canonical ([2]).

2.1.3 The Bratteli diagrams

This graphical presentation of the canonical homomorphism is called a *Bratteli diagram*. Every block of such a diagram is a bipartite graph with the $r \times s$ matrix $\Phi = (\phi_{kl})$. In general, the Bratteli diagram is given by the vertex set V and the edge set E , such that V is an infinite disjoint union $V_1 \sqcup V_2 \sqcup \dots$, where each V_i has a cardinality n . Any pair V_{i-1}, V_i defines a non-empty set $E_i \subset E$ of the edges with a pair of the range and the source functions r, s , such that $r(E_i) \subseteq V_i$ and $s(E_i) \subseteq V_{i-1}$. The non-negative integral matrix of the incidences (ϕ_{ij}) shows how many edges are drawn between the k -th vertex in the row V_{i-1} and l -th vertex in the row V_i . A Bratteli diagram is called *stationary*, if (ϕ_{kl}) is a constant matrix for all $i = 1, \dots, \infty$.

2.2 The number fields

Let $\mathbb{Q}$ be the field of the rational numbers. Let $\alpha \notin \mathbb{Q}$ be an algebraic number over $\mathbb{Q}$, i.e. root of the polynomial equation $a_n x^n + a_{n-1} x^{n-1} + \dots + a_0 = 0$, $a_n \neq 0$, where $a_i \in \mathbb{Q}$. An algebraic extension of the degree n is a minimal field $K = K(\alpha)$, which contains both $\mathbb{Q}$ and α . Note that the coefficients a_i can be assumed integer. If K is an algebraic extension of the degree n over $\mathbb{Q}$, then K is isomorphic to the n -dimensional vector space (over $\mathbb{Q}$) with the basis vectors $\{1, \alpha, \dots, \alpha^{n-1}\}$ (Pollard [8]).

2.2.1 The ring of integers

Let K be an algebraic extension of the degree n over $\mathbb{Q}$. The element $\tau \in K$ is called *algebraic integer* if there exists a monic polynomial $\tau^n + a_{n-1} \tau^{n-1} + \dots + a_0 = 0$, where $a_i \in \mathbb{Z}$. It can be easily verified that the sum and the product of two algebraic integers is an algebraic integer. The (commutative) ring $O_K \subset K$ is called a *ring of integers*. The elements of the subring $\mathbb{Z} \subset O_K$ are called the rational integers. An *integral basis* is a collection $\omega_1, \dots, \omega_n$ of the elements of O_K , whose linear span over the rational integers is equal to O_K .

An integral basis exists for any finite extension and therefore O_K is isomorphic to the integral lattice $\mathbb{Z}^n$, where n is the degree of the field K (Weyl [13]).

2.2.2 The trace of an algebraic number

Let K be a number field of the degree n over $\mathbb{Q}$. There exists n isomorphic embeddings (monomorphisms) $K \rightarrow \mathbb{C}$ (McCarthy [7]). We denote them by $\sigma_1, \dots, \sigma_n$. If $\alpha \in K$ then one defines a *trace* by the formula¹: $\mathcal{N}(\alpha) = \sigma_1(\alpha) + \dots + \sigma_n(\alpha)$. When α is an algebraic integer, then $\mathcal{N}(\alpha)$ is a rational integer. If $p, q \in \mathbb{Z}$, then $\mathcal{N}(p\alpha + q\beta) = p\mathcal{N}(\alpha) + q\mathcal{N}(\beta)$, for all $\alpha, \beta \in K$. It is not hard to see that the above formula establishes a homomorphism $\mathcal{N} : O_K \cong \mathbb{Z}^n \rightarrow \mathbb{Z}$, which does not depend on the choice of an integral basis in O_K (McCarthy [7], Weyl [13]).

2.3 The 3-manifolds

Let M be a compact oriented 3-manifold. Suppose that the second homology group $H_2(M; \mathbb{Z})$ is non-trivial. There exists a linear mapping of the group into the set of the positive integers, which is given by the following construction of W. P. Thurston ([10]).

2.3.1 The Thurston norm

Let X be a connected surface of the genus $g \geq 0$. Denote by $\chi(X)$ its Euler characteristic, i.e. an integer number $2 - 2g$. The negative part of $\chi(X)$ is defined as $\chi_-(X) = \max\{0, -\chi(X)\}$. If X is not connected, one introduces $\chi_-(X)$ as the sum of the negative parts of the connected components of X . For a cycle $z \in H_2(M; \mathbb{Z})$, consider a non-negative integer $N(z) = \inf\{\chi_-(X) \mid X \text{ is an embedded surface representing } z\}$. The $N(z)$ is called a *Thurston norm*. Given two such cycles z_1 and z_2 , let X_1 and X_2 be the surfaces representing them. There exists a unique way to mend X_1 and X_2 together to obtain new embedded surface X , such that $\chi_-(X) = \chi_-(X_1) + \chi_-(X_2)$ (Thurston [10]). Thus, $N(z) = N(z_1) + N(z_2)$ extends linearly to the entire group $H_2(M; \mathbb{Z})$.

2.3.2 The pseudo-Anosov diffeomorphisms

Let X be a surface of the genus $g \geq 2$. Denote by $\text{Mod } X = \text{Diff } X / \text{Diff}_0 X$ the mapping class group of X , i.e. the group of the isotopy classes of the orientation preserving diffeomorphisms of X . The following classification of $\text{Mod } X$ is due to J. Nielsen and W. P. Thurston.

Lemma 1 ([11]) *Any diffeomorphism $\varphi \in \text{Mod } X$ is isotopic to a diffeomorphism φ' , such that either (i) φ' has finite order, or (ii) φ' is pseudo-Anosov*

¹This notion of $\mathcal{N}$ is equivalent to given in section 1.

(non-periodic) diffeomorphism, or (iii) φ' is reducible by a system of curves Γ surrounded by small tubular neighborhoods $N(\Gamma)$, such that on $M \setminus N(\Gamma)$ φ' satisfies either (i) or (ii).

2.3.3 The singularity data

Let φ be a pseudo-Anosov diffeomorphism. There exists a pair of φ -invariant measured foliations $\mathcal{F}_\pm$ on X , such that φ expands along $\mathcal{F}_+$ and contracts along $\mathcal{F}_-$ with dilatation factor $\mu > 1$ (Thurston [11]). $\mathcal{F}_+$ and $\mathcal{F}$ are mutually transversal and have common set of singular points, which are saddle points with $n \geq 3$ prongs. For brevity, we let $\mathcal{F} = \mathcal{F}_+$. Recall that the index of n -prong saddle s_n is $-\frac{1}{2}(n-2)$. Therefore $\sum_{s_n \in \text{Sing } \mathcal{F}} \frac{n-2}{2} = 2g-2$, where g is the genus of surface X . If $m = |\text{Sing } \mathcal{F}|$ is the total number of the singular points of $\mathcal{F}$, then $1 \leq m \leq 4g-4$, where the minimum is attained by a unique saddle s_{4g-2} and maximum by the set $\{s_3, s_3, \dots, s_3\}$ of $4g-4$ saddles. We refer to the set $\{s_{i_1}, \dots, s_{i_m}\}$ as a *singularity data* of $\mathcal{F}$.

2.3.4 The mapping tori

Let $\varphi : X \rightarrow X$ be a diffeomorphism of the surface X . One can obtain 3-dimensional manifolds $M = M(\varphi)$ by the formula $M = \{X \times [0, 1] \mid (x, 0) \mapsto (\varphi(x), 1), x \in X\}$. The manifold M is called a *mapping torus*. It is not hard to see that M is a mapping torus if and only if $M \rightarrow S^1$ is a fibre bundle over S^1 with the monodromy φ . If the diffeomorphism $\varphi \in \text{Mod } X$ is of a finite order, then M will be a Seifert manifold. In the case when φ is pseudo-Anosov, the following result due to W. P. Thurston is true.

Lemma 2 ([12]) *The mapping torus M admits a hyperbolic structure, if and only if the diffeomorphism φ is pseudo-Anosov.*

2.3.5 The second homology group of the mapping torus

Let $\varphi : X \rightarrow X$ be a pseudo-Anosov diffeomorphism of genus $g \geq 2$ surface. Let $\text{Sing } \mathcal{F}$ be a finite set of the singularities of the φ -invariant foliation $\mathcal{F}$. The relative homology group $H_1(X, \text{Sing } \mathcal{F}; \mathbb{Z})$ is a torsion-free of the rank $k = 2g + |\text{Sing } \mathcal{F}| - 1$, where $|\text{Sing } \mathcal{F}|$ is the cardinality of the set $\text{Sing } \mathcal{F}$. Let M be the mapping torus of φ . The 2-cycles of M are generated by the 1-cycles of $X \setminus \text{Sing } \mathcal{F}$. Indeed, let $C \in H_1(X, \text{Sing } \mathcal{F}; \mathbb{Z})$ and $\varphi(C)$ its image under the diffeomorphism φ . Let $X - C$ be a copy of the surface X , which is cut along the closed curve C . Similarly, let $X - \varphi(C)$ be a copy of X with a cut along $\varphi(C)$. The surface $X = (X - C) \cup (X - \varphi(C))$, glued along the action of φ , belongs to the group $H_2(M)$, and any non-torsion element of $H_2(M)$ can be obtained in such a way. Therefore, $\text{rank } H_2(M) = 2g + m - 1$, where $m = |\text{Sing } \mathcal{F}|$.

2.3.6 The Thurston norm of the surface bundle

It is interesting to relate the Thurston norm N on the second homology with the geometry of M . It turns out, that in this case N can be expressed in terms of the Euler classes of the plane bundle tangent to the fibres of M . Namely, let N^* be a dual Thurston norm defined on the first homology $H_1(M; \mathbb{Z})$ by the formula $N^*(z) = \sup_{N(u) \leq 1} u(z)$, where $u \in \text{Hom}(H_1(M; \mathbb{Z}), \mathbb{Z}) \simeq H^1(M; \mathbb{Z})$ and N is the Thurston norm on $H^1(M; \mathbb{Z}) \simeq H_2(M; \mathbb{Z})$ induced by the Poincaré duality. Then the following lemma is true.

Lemma 3 ([3]), ([10]) *Let τ be a subbundle of the tangent bundle TM consisting of the 2-planes tangent to the fiber X of the fibration $M \rightarrow S^1$. Let $e(\tau) \in H^2(M; \mathbb{Z})$ be the Euler class of τ , i.e. first obstruction to the cross-section of bundle τ . Then (i) the norm $N^* : H_1(M; \mathbb{Z}) \rightarrow \mathbb{Z}^+$ is induced by the cocycle $e(\tau)$, i.e. $N^*(z) = |\int_z e(\tau)|$; (ii) the set of the (de Rham) cohomology classes $H^1(M; \mathbb{R})$, which is representable by the closed non-singular differential 1-forms on M is a maximal cone $C \subset H^1(M; \mathbb{R})$, where the Thurston norm N can be extended linearly.*

3 Proof of theorem 1

(i) Let $\mathcal{N} : H_2(M) \rightarrow \mathbb{Z}$ be a linear mapping. To show that the set $C = \{z \in H_2(M) \mid \mathcal{N}(z) > 0\}$ is a cone, we have to establish that

- (1) $z \in C, c > 0$ implies $cz \in C$
- (2) $z_1, z_2 \in C$ implies $z_1 + z_2 \in C$.

Indeed, since $\mathcal{N}$ is linear, $\mathcal{N}(cz) = c\mathcal{N}(z)$, where $\mathcal{N}(z) > 0$ by the assumption. Therefore, $c\mathcal{N}(z) > 0$ and the item (1) follows. Similarly, in the item (2), by the linearity of $\mathcal{N}$, we have $\mathcal{N}(z_1 + z_2) = \mathcal{N}(z_1) + \mathcal{N}(z_2) > 0$, since $\mathcal{N}(z_1) > 0, \mathcal{N}(z_2) > 0$ by the assumption. The item (i) is proved.

(ii) The proof of item (ii) is based on a lemma of D. Gabai ([4]). Roughly speaking, we shall estimate the Gromov norm (a simplicial norm) of $H_2(M)$, rather than the Thurston norm itself. This approach gives a technical advantage, because the group of the 2-chains in M has a natural abelian structure. Next we use the Gabai lemma to evaluate the two norms. The Gromov norm was introduced and studied in ([5]).

Let M be a compact manifold and $z \in H_2(M)$ be an element of the second homology group of M . A *Gromov norm* $g(z)$ is given by the formula $g(z) = \inf_{Z \in [z]} \{\sum |a_i| : Z = \sum a_i \sigma_i\}$, where $[z]$ is the homology class of the 2-chains and $\sigma_1, \dots, \sigma_n$ is a basis of the simplicial decomposition of M . The following lemma is true.

Lemma 4 *Suppose M is a compact oriented 3-manifold. Then the Thurston norm N and Gromov norm g are related by the formula $N(z) = \frac{1}{2}g(z)$, for each $z \in H_2(M)$ in the domain of definition of the two norms.*

Proof. The proof can be found in ([4]), Corollary 6.18. For the sake of clarity, let us outline the main idea. First, notice that if M is a hyperbolic k -manifold and $[M]$ its homology class, then we have Gromov's formula $g([M]) = \text{Vol } M / \text{Vol } \sigma$, where σ is the largest hyperbolic k -simplex (Gromov [5]). Thus, for the connected surface X , one has $g([X]) = \frac{2\pi|\chi(X)|}{\pi} = 2N([X])$. The formula extends to the case X with more than one connected component and requires the singular norms x_s in this case, see Gabai ([4]). Eventually, it can be shown, that $x_s = N$ and one gets inequality $g \leq 2N$.

To prove the inequality $g \geq 2N$, let $z \in H_2(M)$ and $Z \in [z]$ be a 2-cycle $Z = \sum a_i \sigma_i$, where $a_i \in \mathbb{Z}$. By pasting the singular simplices, one can obtain a proper map $f : X \rightarrow M$, such that $[f(X)] = z$. By Gromov's formula for the hyperbolic volumes $2N([X]) \leq \sum |a_i| = g([X])$. Lemma 4 follows. $\square$

Fix a simplicial basis $\sigma_1, \dots, \sigma_n$ in the group $\mathcal{C}$ of the 2-chains of the regular triangulation of M . Consider a subset of $\mathcal{C}$, given by the formula $K_{\mathcal{C}} = \{Z \in \mathcal{C} \mid Z = \sum_{i=1}^n a_i \sigma_i, \quad a_i > 0\}$. Denote by $\tilde{\mathcal{N}} : H_2(M) \rightarrow \mathbb{Z}^+$ the mapping $z \mapsto g(z)$ given by the Gromov norm. It is not hard to verify, that $\tilde{\mathcal{N}}$ is linear on the $K_{\mathcal{C}}$ and the $K_{\mathcal{C}}$ is a cone in $\mathcal{C}$. Consider the following commutative diagram of the linear mappings:

$$\begin{array}{ccc} K_{\mathcal{C}} & \xrightarrow{\Sigma} & \mathcal{C} \\ \tilde{\mathcal{N}} \downarrow & & \downarrow \mathcal{N} \\ \mathbb{Z}^+ & \xrightarrow{w} & \mathbb{Z}^+ \end{array}$$

(Here w is a doubling map, acting by the formula $z \mapsto 2z$). The map Σ on the diagram is linear. The Σ establishes a bijection between the bases $\{\sigma_i\}$ and $\{\omega_i\}$: $\Sigma(\sigma_i) = \omega_i$. It remains to apply lemma 4. The item (ii) of theorem 1 follows. $\square$

Acknowledgments. I wish to thank O. Bratteli, D. Calegari, M. Felisatti, Y. Minsky and V. Turaev for helpful comments.

References

- [1] O. Bratteli, P. E. T. Jørgensen, K. H. Kim, and F. Roush, Computation of isomorphism invariants for stationary dimension groups. *Ergodic Theory Dynam. Systems* 22 (2002), 99-127.

- [2] E. G. Effros, Dimensions and C^* -Algebras, in: Conf. Board of the Math. Sciences No.46, AMS (1981).
- [3] D. Fried, Fibrations over S^1 with pseudo-Anosov monodromy, in: A. Fathi, F. Laudenbach and V. Poénaru, Travaux de Thurston sur les Surfaces, Exposé 14, Astérisque 66-67 (1979), 251-266.
- [4] D. Gabai, Foliations and the topology of 3-manifolds, J. Differential Geometry 18 (1983), 445-503.
- [5] M. Gromov, Volume and bounded cohomology, Inst. Hautes Études Sci. Publ. Math. 56 (1982), 5-99.
- [6] D. Handelman, Positive matrices and dimension groups affiliated to C^* -algebras and topological Markov chains, J. Operator Theory 6 (1981), 55-74.
- [7] P. J. McCarthy, Algebraic Extensions of Fields, Blaisdell Publishing Company, 1966.
- [8] H. Pollard, The Theory of Algebraic Numbers, Carus mathematical monographs, 9.
- [9] M. Rørdam, F. Larsen and N. Laustsen, An introduction to K -theory for C^* -algebras. London Mathematical Society Student Texts, 49. Cambridge University Press, Cambridge, 2000. xii+242 pp. ISBN: 0-521-78334-8; 0-521-78944-3
- [10] W. P. Thurston, A norm for the homology of 3-manifolds, Mem. Amer. Math. Soc. 339 (1986), 99-130.
- [11] W. P. Thurston, On the geometry and dynamics of diffeomorphisms of surfaces, Bull. Amer Math. Soc. 19 (1988), 417-431.
- [12] W. P. Thurston, Hyperbolic structures on 3-manifolds, II: surface groups and 3-manifolds which fiber over the circle, arXiv:math.GT/9801045 v1.
- [13] H. Weyl, Algebraic Theory of Numbers, Annals of Math. Studies 1, Princeton Univ. Press, 1940.

THE FIELDS INSTITUTE FOR MATHEMATICAL SCIENCES, TORONTO, ON,
CANADA, E-MAIL: igor.v.nikolaev@gmail.com

Current address: 101-315 Holmwood Ave., Ottawa, ON, Canada, K1S 2R2